arXiv:1504.00660v1 [math.SP] 1 Apr 2015

A BOUND FOR RATIOS OF EIGENVALUES OF SCHRÖDINGER OPERATORS WITH SINGLE-BARRIER POTENTIALS.

JAMEL BEN AMARA AND HEDHLY JIHED

ABSTRACT. M. Horvath and M. Kiss [Proc. Amer. Math. Soc, (2005)] , proved that the upper estimate $\frac{\lambda_n}{\lambda_m} \leq \frac{n^2}{m^2}$ $(n > m \geq 1)$ of Dirichlet Shrodinger operators with nonnegative and single-well potentials. In this paper we discuss the case of nonpositive potentials $q(x)$ defined and continuous on the interval $[0,1]$. Namely, we prove that if $q(x) \leq 0$ and single-barrier then $\frac{\lambda_n}{\lambda_m} \geq \frac{n^2}{m^2}$ for $\lambda_n > \lambda_m \geq -2q_-$, where $q_- = \min\{q(x),\ x \in [0,1]\}$. Moreover we show that there exixst $l_0 \in (0,1]$ such that if $q(x) \leq 0$ and single-barrier, then the associated eigenvalues $(\lambda_n(\ell_0))_{n\geq 1}$ satisfy $\lambda_1(\ell_0) > 0$ and $\frac{\lambda_n(\ell_0)}{\lambda_m(\ell_0)} \geq \frac{n^2}{m^2}$ for $n > m \geq 1$.

1. INTRODUCTION

Consider the Shrodinger operator

$$-y'' + q(x)y = \lambda y, \quad x \in [0,1] \tag{1.1}$$

where $q \in C[0,1]$, with Dirichlet boundary conditions

$$y(0) = y(1) = 0. \tag{1.2}$$

It is known (see [6]) that the spectrum of problems $(1.1)-(1.2)$ consists of a growing sequence of infinitely point $\lambda_1 < \lambda_2 < .... < \lambda_n ...$.

Ashbaugh and Benguria [1] proved an optimal upper estimate of the eigenvalue ratio

$$\frac{\lambda_n}{\lambda_1} \leq n^2$$

of Dirichlet Shrodinger operator with nonnegative potentials $q(x)$. They also examined the ratio of two arbitrary eigenvalues, and found

$$\frac{\lambda_n}{\lambda_m} \leq \left\lceil \frac{n}{m} \right\rceil^2 \text{ for } m < n,$$

where $\lceil s \rceil$ denotes the smallest integer greater than or equal to s. Yu-Ling Huang and C. K. Law [3] extented the results of [1] to the cases of Equation (1.1) with $q(x) \geq 0$ and general boundary conditions. Chung-Chuan Chen [4] proved the bound

$$\frac{\lambda_n}{\lambda_m} \geq \left(\left\lfloor \frac{n}{m} \right\rfloor \right)^2$$

for problem $(1.1) - (1.2)$ with $q(x) \leq 0$, where $\lfloor s \rfloor$ denotes the smallest integer less than or equal to s. M. Horváth and M. Kiss [5] showed that for nonnegative single-well potentials,

$$\frac{\lambda_n}{\lambda_m} \leq \frac{n^2}{m^2}.$$

Let $0 \leq x_0 \leq 1$ *be fixed. Following Ashbaugh and Benguria* [2] *we call the function* q *a single-well function if* q *is decreasing in* $[0, x_0]$ *and increasing in* $[x_0, 1]$. *Analogously,* q *is called a single-barrier function if it is increasing in* $[0, x_0]$ *and decreasing in* $[x_0, 1]$.

In this paper we investigate the case of nonpositive potentials $q(x)$ defined and continuous on the interval $[0,1]$. Namely, we prove that if $q(x) \leq 0$ and single-barrier then $\frac{\lambda_n}{\lambda_m} \geq \frac{n^2}{m^2}$ for $\lambda_n > \lambda_m \geq -2q_{\text{-}}$, where $q_{\text{-}} = \min\{q(x),\ x \in [0,1]\}$. Moreover we show that there exixst $l_0 \in (0,1]$ such that if $q(x) \leq 0$ and single-barrier, then the associated eigenvalues $(\lambda_n(\ell_0))_{n\geq 1}$ (of Problem (1.1) – (1.2) defined on $[0, \ell_0]$) satisfy $\lambda_1(\ell_0) > 0$ and $\frac{\lambda_n(\ell_0)}{\lambda_m(\ell_0)} \geq \frac{n^2}{m^2}$ for $n > m \geq 1$.

2. PRELIMINARIES AND THE MAIN STATEMENTS

In this section, we introduce the modified Prüfer transformation that is the key to our results.

Denote by $y(x,z)$ the unique solution of the initial value problem

$$-y'' + q(x)y = z^2 y,\ x \in [0,1],\ z > 0 \tag{2.1}$$

$$y(0) = 0,\ y'(0) = 1 \tag{2.2}$$

The Prüfer variable $r(x,z)$, $\varphi(x,z)$ that we use here are defined by

$$y(x,z) = r(x,z)\sin\varphi(x,z) \tag{2.3}$$

$$y'(x,z) = zr(x,z)\cos\varphi(x,z) \tag{2.4}$$

$$\varphi(0,z) = 0, \tag{2.5}$$

where $r(x,z) > 0$, and we denote by prime the derivative with respect to x (and by dot the derivative with respect to z). Define further

$$\theta = \frac{\varphi}{z} \tag{2.6}$$

Using Equation (1.1) one finds the following differential equations for $r(x,z)$, $\varphi(x,z)$:

$$\varphi' = z - \frac{q}{z}\sin^2\varphi \tag{2.7}$$

$$\frac{r'}{r} = \frac{q}{z}\sin\varphi\cos\varphi. \tag{2.8}$$

It is obvious that $y = 0$ iff $\sin\varphi = 0$, hence z^2 is an eigenvalue iff $\varphi(\pi, z)$ is a multiple of π. Denote by z_n the square root of λ_n.

Theorem 1. *Let* $q(x) \leq 0$ *be monotone increasing in* $[0, x_0]$. *Then* $\dot{\theta}(x_0, z) \leq 0$ *for* $z \geq \sqrt{-2q_-}$, *where* $q_- = \min\{q(x),\ x \in [0,1]\}$. *If there is a* $z \geq \sqrt{-2q_-}$ *with* $\dot{\theta}(x_0, z) = 0$, *then* $q = 0$ *in* $(0, x_0]$.

Theorem 2. *For the regular Sturm-Liouville system* $(1.1) - (1.2)$, *if* $q(x) \leq 0$, *continuous and single-barrier, then the* n^{th} *and* m^{th} *eigenvalues with* $\lambda_n > \lambda_m \geq -2q_-$ *(where* $q_- = \min\{q(x),\ x \in [0,1]\}$*),*

$$\frac{\lambda_n}{\lambda_m} \geq \frac{n^2}{m^2}.$$

and if for two different m *and* n *equality holds if and only if* $q = 0$ *in* $[0,1]$.

Theorem 3. *There exist* $\ell_0 \in (0,1]$ *such that if* $q \leq 0$, *continuous and single-barrier, then* $\lambda_1(\ell_0) > 0$ *and for the* n^{th} *and* m^{th} *eigenvalues with* $n > m \geq 1$, *(of Sturm-Liouville Problems* $(1.1) - (1.2)$ *defined on* $[0, \ell_0]$*)*

$$\frac{\lambda_n(\ell_0)}{\lambda_m(\ell_0)} \geq \frac{n^2}{m^2},$$

and if for two different m *and* n *equality holds if and only if* $q = 0$ *in* $[0, \ell_0]$.

Theorem 4. *Consider Equation* (1.1) *with the Dirichlet-Neumann boundary conditions*

$$y(0) = y'(1) = 0. \tag{2.9}$$

If the potential $q(x) \leq 0$, *continuous and increasing, then for the* m^{th} *and* n^{th} *eigenvalues with* $\lambda_n > \lambda_m \geq -2q_-$, *(where* $q_- = \min\{q(x),\ x \in [0,1]\}$*),*

$$\frac{\lambda_n}{\lambda_m} \geq \frac{(2n-1)^2}{(2m-1)^2},$$

and if for two different m *and* n *equality holds, then* $q = 0$ *in* $(0,1]$.

References

[1] M. S. Ashbaugh and R. D. Benguria, Optimal bounds for ratios of eigenvalues of one dimensional Schrödinger operators with Dirichlet boundary conditions and positive potentials, Comm. Math. Phys. 124 (1989)

[2] M. S. Ashbaugh and R. D. Benguria, Optimal lower bound for the gap between the rst two eigenvalues of one-dimensional Schrodinger operators with symmetric single-well potentials, Proc. Amer. Math. Soc. 105, 419 – 424 (1989).

[3] Yu-Ling Huang and C. K. Law, Eigenvalue ratios for the regular Sturm-Liouville system. Proceedings of the American Mathematical Society. 124, (1996) .

[4] Chung-Chuan Chen, Optimal lower estimates for eigenvalue ratios of Schrödinger operators and vibrating strings, electronically published e-Thesis (2002).

[5] Miklós Horváth and Márton Kiss, A bound for ratios of eigenvalues of Schrödinger operators with single-well potentials. Proceedings of the American Mathematical Society. 134, (2005) .

[6] B. M. Levitan and I. S. Sargsjan, Sturm-Liouville and Dirac operators (in Russian), Nauka, Moscow, (1988).

Faculté des Sciences de Tunis, Université El Manar de Tunis
E-mail address: `jamel.benamara@fsb.rnu.tn`

Faculté des Sciences de Tunis, Université El-Manar de Tunis
E-mail address: `hjihed@gmail.com`